

AD741702

Best Available Copy

Reproduced by
**NATIONAL TECHNICAL
INFORMATION SERVICES**
Springfield, Va. 22161

TWO ITEMS CONCERNING DIRECTIONAL DATA

BY

M. A. STEPHENS

TECHNICAL REPORT NO. 192

April 3, 1972

PREPARED UNDER CONTRACT N00014-67-A-0112-0053


(NR-042-267)

OFFICE OF NAVAL RESEARCH

Herbert Solomon, Project Director

Reproduction in Whole or in Part is Permitted for
any Purpose of the United States Government

DEPARTMENT OF STATISTICS
STANFORD UNIVERSITY
STANFORD, CALIFORNIA



UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the original report is classified)

1. ORIGINATING ACTIVITY (Corporate outside)		2A. REPORT SECURITY CLASSIFICATION	
Department of Statistics Stanford University			
3. REPORT TITLE		2B. GROUP	
TWO ITEMS CONCERNING DIRECTIONAL DATA			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
TECHNICAL REPORT			
5. AUTHOR(5) (Last name, first name, initial)			
STEPHENS, M. A.			
6. REPORT DATE	7B. TOTAL NO. OF PAGES	7A. NO. OF REFS	
April 3, 1972	16	6	
8A. CONTRACT OR GRANT NO.	9A. ORIGINATOR'S REPORT NUMBER(5)		
N00014-67-A-0112-0053	Technical Report No. 192		
9. PROJECT NO.	9B. OTHER REPORT NO(5) (Any other numbers that may be assigned this report)		
NR-042-267			
10. AVAILABILITY/LIMITATION NOTES			
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Statistics & Probability Program Office of Naval Research Arlington, VA.	
13. ABSTRACT.			
<p>1. A discussion is given of confidence procedures for the model vector \underline{A} of a von Mises or Fisher distribution. Slight revisions of confidence levels are suggested because the usual procedure strictly gives the axis but not the direction of \underline{A}.</p> <p>2. The likelihood ratio test is given, with percentage points, for testing $\underline{A} = \underline{A}_0$, for the Fisher distribution. A good approximation is offered for the similar test for the von Mises distribution.</p>			

DD FORM 1473
1 JAN 64

UNCLASSIFIED

Security Classification

UNCLASSIFIED
Security Classification

16. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
directional data						
vectors						
Fisher distribution						
von Mises distribution						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).
10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

TWO ITEMS CONCERNING DIRECTIONAL DATA

by

M.A. Stephens

In this paper we consider two topics connected with directional data, i.e., observations which may be recorded by unit vectors OP_1 from the center O of a circle or sphere of radius 1, to points P_1 on the circumference or surface. Alternatively, the point P_1 on a circle may record an observation, such as the occurrence of an event during a period whose length is represented by the total circumference.

When the observations are clustered around a central direction, the von Mises distribution, on the circle, or the Fisher distribution, on the sphere, are used to describe the data. These are unimodal probability distributions, with density on the surface proportional to $\exp(k\cos\alpha)$, where k is a concentration parameter and α the angle between OP and the central (modal) vector A . It will be assumed that the reader is familiar with these distributions; the references given in the text may be used as sources for earlier references.

The two topics to be discussed in the next two sections are. In section 2; A discussion of the standard tests and confidence interval procedures for the modal vector A of a von Mises or a Fisher distribution, when the concentration parameter k is not known. It is not generally noted that the usual test is strictly for the axis along which A lies, without specifying the direction, and this necessitates a slight revision of confidence levels.

In section 3, Significance points are provided for the likelihood-ratio test for \underline{A} for a Fisher distribution, for the case when the concentration parameter κ is known. The test will replace a test already suggested elsewhere, and indications are that an approximation will hold well for the von Mises distribution also.

2. Tests and Confidence Intervals for the Modal Vector of a von Mises or Fisher Distribution.

2.1. The von Mises and Fisher distributions have densities proportional to $\exp(\kappa \cos \alpha)$, where α is the angle between a sample vector \underline{OP} and the modal vector \underline{A} , and κ is a concentration parameter. For a test of H_0 : \underline{A} is along a given \underline{A}_0 , when κ is not known, Watson and Williams (1956) proposed a conditional test, both for the circle and the sphere. Suppose a sample of N unit vectors gives a resultant vector \underline{R} , length R , and let its component on \underline{A}_0 be X ; the test depends on the fact that the conditional distribution of R , given X , is independent of κ . Critical values R_0 , for given N , X , and α , have been given by Stephens (1962a,b) in the form of charts of R_0 against X ; if R exceeds R_0 , the null hypothesis is rejected.

The distribution of R , given X , is in fact the same for X positive or negative, and if a given \underline{A}_0 is acceptable when the component of \underline{R} on \underline{A}_0 is positive, the vector $-\underline{A}_0$ would be equally acceptable, the component being now negative. Thus the test is strictly a test for the modal axis, without direction. This will become important when we use critical values R_0 of R to obtain a confidence interval for \underline{A} . The use of the charts both for testing and for confidence

intervals, can best be illustrated with the help of a diagram. In Figure 1 we have taken the case of a sample of size $N = 20$, drawn from the von Mises distribution.

Suppose OX points in the direction of the modal vector A , and let OY be the axis at right angles. For every sample with resultant R , we can calculate $Y = \pm \sqrt{R^2 - X^2}$, and set $C \equiv (X, Y)$. OX is the resultant R , and C represents the sample on the diagram. C lies inside the circle, center O , radius 20. Let R_0 be the critical value of R for given X , at the 5% level, given by the charts, and draw the curve $Y^2 = R_0^2 - X^2$; call this the 5% critical limit. This is shown in the illustration, and also part of the 1% critical limit obtained from 1% critical values of R . The curves are symmetrical about OY as well as OX . Since for every X the probability of C falling outside the curve shown is 0.05, with Y either positive or negative, the probability of falling above the upper curve is 0.025 and of falling below the lower curve is also 0.025, whatever the value of κ . Suppose now OX has coordinate θ_0 measured anticlockwise from a suitable initial line OB , if the true modal vector were along OX' with angle $\theta_1 > \theta_0$, more than 2.5% of sample points C would fall above the upper 2.5% curve. In the same way, if $\theta_1 < \theta_0$ an excess of sample points will lie below the lower 2.5% critical boundary. As a result, a test of the hypothesis, H_0 , that the direction of the modal vector is $\theta_c = \theta_0$, against the one-sided alternative that $\theta_c > \theta_0$, consists in (a) Calculating X and R from the sample, (b) finding from the charts in Stephens (1962a) the critical value R_0 of $R(.05)|X$ (or $R(.01|X)$) and (c) rejecting H_0 at the

2.5%(or 0.5%) level, if the observed $R > R_0$. There will be a corresponding one-sided test against the alternative $\theta_c < \theta_0$, and if the alternative is simply $\theta_c \neq \theta_0$, then the steps above are followed but, in (c), H_0 is rejected, if R is too large, at the 5% (or 1%) level. For N, X not given in the charts, Stephens (1962a) has given several approximations. Similar charts and approximations for use with the Fisher distribution, are given in Stephens (1962b).

2.2 Confidence limits for θ_c .

Suppose that the true modal vector has direction $\theta_c = \theta_0$, but that we are unaware of this and must estimate θ_c from the direction of the sample vector resultant. Figure 1 shows two possible resultants OC_1, OC_2 , at angles θ_1, θ_2 respectively, both of the same length $R = 7.90$. We find from Stephens' tables that for $N = 20$ the corresponding abscissa for $\alpha = 0.05$, is $X = 5.0$, and note that

$$\phi = \cos^{-1}(X/R) = \cos^{-1} = \cos^{-1}(5.0/7.9) = \cos^{-1}(.6329) = 50.7^\circ.$$

ϕ is clearly the angle between OZ and OX , where OZ is 7.9 and Z is on the 5% critical limit.

Suppose the lower confidence limit for θ_c is obtained by subtracting ϕ from the angle of R ; the two illustrations then give limits $\theta_1 - 50.7^\circ$ and $\theta_2 - 50.7^\circ$. In the first example this limit gives a confidence interval which does not include the true modal vector, along OX , and this is clearly because the point C_1 is above the 5% critical boundary, while the interval based on OC_2 does include OX , because C_2 is below the 5% boundary. Since 2.5% of samples

would give a sample point above the boundary, like C_1 , and 97.5% below, like C_2 , the procedure clearly provides a lower 97.5% confidence limit for θ_c , whatever is the value of κ . Similarly, if we add $\phi = 50.7^\circ$ to the angle of the sample resultant vector we shall have an upper 97.5% confidence limit for θ_c , and the two limits will define a central 95% confidence interval for θ_c . A similar result holds for the left side of the diagram, where X is negative. Thus the interval is not strictly for the modal vector with direction, but only for the axis along which it lies, with either possible direction. In practice, of course, except for κ very small, negative values of X will rarely occur, and the confidence interval is always chosen to give positive X , but this lowers the confidence level. The amount by which it is lowered is found as follows. Let p be the probability, for given κ , that $X > 0$, i.e., the probability that the sample point falls to the right of OY . If P is the confidence probability (i.e., 100% is the confidence level) then the probability that the above procedure includes the positive modal vector is Pp . The relationship of p and κ , for p near 1, may be found approximately from a table of percentage points of X , in Stephens (1969a). From this we have

N:	10	10	20	20	40	40
p:	0.95	0.99	0.95	0.99	0.95	0.99
κ :	0.74	1.08	0.53	0.74	0.37	0.53

Thus for $\kappa > 1$, the risk of mistaking the direction of \hat{A} will be very small, for $N = 10$ say, and the confidence level hardly changes,

for larger samples, κ may become even smaller with negligible risk of a wrong decision.

2.3 The Fisher Distribution.

For three-dimensions, the same type of argument holds, though a geometric representation would now involve an ellipsoid for the 5% critical limit. Corresponding to two-sided confidence intervals for θ on the circle, there will now be a cone of confidence for θ on the sphere; it would be difficult to interpret one-sided tests or confidence intervals. Again the test based on R for given X is really a test for the axis of A , without direction, and the alleged confidence probability P must be multiplied by a p which depends on N and κ . From tables of the distribution of X for given N , κ given in Stephens (1967) the table of values for p for the sphere becomes

N:	10	10	20	20	40	40
p:	0.95	0.99	0.95	0.99	0.95	0.99
κ :	0.91	1.30	0.64	0.91	0.45	0.64

κ is required to be a little higher for the sphere than for the circle to obtain the same p for given N .

2.4 Numerical illustration.

In Table 1 we give, for $N = 10$ and 20 , the relationship between R/N and the critical value X/N for a 95% confidence interval with the Fisher distribution, also the θ_c to which this corresponds.

Further, R/N is used to estimate κ (by $\hat{\kappa}$, the solution of $\coth \kappa - 1 - \kappa = R/N$); the estimate could then be used to estimate p as described above. Values of $\hat{\kappa}$ are also included in Table 1.




Fig 1. $N=20$

TABLE 1

Critical values of X/N and θ_c (to nearest degree) for 95% confidence interval, given R/N , and estimated value $\hat{\kappa}$ derived from R/N . (Fisher distribution).

N = 10								
R/N:	.44	.45	.5	.55	.6	.7	.8	.9
X/N:	.00	.056	.241	.342	.431	.582	.721	.861
θ_c :	90	83	61	52	44	34	26	17
$\hat{\kappa}$:	1.51	1.55	1.80	2.07	2.40	3.30	5.00	10.00

N = 20								
R/N:	.32	.35	.4	.5	.6	.7	.8	.9
X/N:	.00	.16	.29	.42	.53	.65	.77	.88
θ_c :	90	63	44	34	28	22	17	11
$\hat{\kappa}$:	1.03	1.14	1.34	1.80	2.40	3.30	5.00	10.00

3. Likelihood-Ratio Test for the Modal Vector of a Fisher Distribution When the Concentration Parameter is Known.

3.1 The test to be discussed in this section is similar to that in Section 2, but with the concentration parameter κ assumed known. This is probably less likely to occur, but when it does, a more powerful test of H_0 can be made. The likelihood-ratio test statistic for $H_0: \underline{A}$ along \underline{A}_0 , against the alternative $H_A: \underline{A}$ is along a vector other than \underline{A}_0 , may be shown to be $R \cdot X$. Thus if the distribution of $Z = R \cdot X$ were known, a test based on Z , rejecting if Z were too large, should give a more powerful test against the alternative. The exact distribution of Z is difficult to find; however, for the Fisher distribution, significance points can be found, and are given in Table 2. The test may formally be set out as follows:

- (a) Calculate $Z = R \cdot X$;
- (b) In Table 2 for given N , κ and α , find the table entry z ;
- (c) If $Z > z$, reject H_0 at significance level α .

If κ is too large for Table 2, solve for z from

$$2\kappa z = \chi^2_2(\alpha), \quad (1)$$

where $\chi^2_2(\alpha)$ is the upper significance point of χ^2_2 at level α .

3.2 Theory of the test.

We start by defining

$$c(\kappa) = \left(\frac{\kappa}{2 \sinh \kappa} \right)^N, \text{ and}$$

$$P_N^r(t) = \sum_{s=0}^{\infty} \binom{N}{s} (-1)^s \langle N-t-2s \rangle^r,$$

N and r are positive integers, and $|t| < N$, the notation $\langle z \rangle$ means $\langle z \rangle = z$ if $z > 0$, and $\langle z \rangle = 0$ if $z \leq 0$. The joint density of R and X is (Stephens, 1967)

$$f_1(R, X) = c(\kappa) \exp(\kappa X) P_N^{N-2}(R) / (N-2)!, \quad |X| < R, \quad 0 < R < N; \quad (2)$$

Let $y = R - X$, the joint density of y, R is then

$$f_2(y, R) = c(\kappa) \exp(-\kappa y) \exp(\kappa R) P_N^{N-2}(R) / (N-2)!;$$

$$0 \leq y \leq 2R, \quad 0 \leq R \leq N \quad (3)$$

For the density of y alone, we must integrate out R , so we require

$$J = \int_v^N \exp(\kappa R) P_N^{N-2}(R) dR, \text{ with } v = y/2.$$

This integral J enters an expression for an integral I given in Stephens (1967, top of page 216) and may easily be deduced from the result for I , the density of y then becomes

$$f_3(y) = \kappa c(\kappa) \exp(-\kappa y) \left[\exp(-\kappa y/2) \sum_{r=2}^{N-1} \frac{P_N^{N-r}(y/2)}{\kappa^r (N-r)!} + \frac{1}{\kappa^N} \sum_{r=0}^t (\exp(\kappa(N-2r)) - \exp(\kappa y/2)) \binom{N}{r} (-1)^r \right], \quad 0 \leq y \leq 2N.$$

Where t is the greatest integer less than $(N-y/2)/2$. This density has been integrated numerically to give the significance points in Table 2.

The χ^2 approximation in step (c) of Section 3.2 may be deduced from the fact that $-2 \ln \lambda = 2\kappa(R-X)$, where λ is the likelihood ratio for the test of H_0 , and so is asymptotically χ^2 distributed. However, Watson (1956) first gave the approximation, and deduced it from the properties of the Fisher density for large κ . It is interesting also to see that it follows from (2) above, if κ is large, since in this circumstance we expect large R and X and therefore small values of y . Thus if we replace the range of y in (2) by $0 \leq y \leq \omega$, and adjust the constant $c(\kappa)$ accordingly, we can see that the joint density of y and R now factors into two components, one containing only y and the other only R ; with appropriate choice of constant terms, these must be the marginal densities of y and R . For y , since the term with y is $\exp(-\kappa y)$, we see that κy must have the exponential distribution, the full density is

$$f(y) = \kappa \exp(-\kappa y), \quad y > 0$$

and this is equivalent to

$$2\kappa y \approx \chi_2^2. \quad (3)$$

These two approximate derivations of (3) show that we can expect it to be a good approximation: (a) when κ is large, for all N , and (b) for smaller values of κ , as $N \rightarrow \infty$. The values given by the approximation are recorded in Table 2 opposite $N = \infty$, the exact values given for other values of κ show that the above expectations are borne out extremely well.

3.3 Power comparisons.

The test of H_0 above can also be made using X alone, since its distribution for given N and κ is known and significance points tabulated. This suggestion was given in Stephens (1967), but was wrongly described as a two-tailed test, in fact, H_0 would be rejected only if X is too small for given N and κ . We now compare the two tests for power, not directly, but indirectly by comparing the confidence intervals for A obtained by the two methods. Suppose A is at angle θ to B ; a $(1-\alpha)\%$ confidence interval for θ is $0 \leq \theta \leq \theta_c$, where θ_c is the solution of $R \cos \theta_c = X_c$, and X_c is the critical value of X found in either test, at significance level α .

Consider an example, when $N = 10$ and $\kappa = 4$. The 5% critical value of $R-X$ is found from step (c), the value is 0.750, so $X_c = R - 0.750$. For the test based on X alone, $X_c = 6.10$ (from Table 1, Stephens (1967)). Thus the $R-X$ test will give smaller confidence band for θ whenever $R - 0.750 > 6.10$, i.e., whenever $R > 6.85 = R_0$ say. From Table 2 of Stephens (1967) the probability $\Pr(R > R_0)$ can roughly be estimated, for $N = 10$, $\kappa = 4$, the lower 1% and 5% values

of R are 6.42 and 5.71 and $\Pr(R > 6.85)$ is near 0.9. Thus 90% of the time the R - X test, for $N = 10$, $\kappa = 4$, would give greater power than the test based on X alone. For $N = 10$, and other values of κ , the value of R_0 is given, and a rough estimate of the probability p of exceeding it, in the following small table:

κ :	1	1.5	2	2.5	3	4
R_0 :	3.21	3.78	4.56	5.29	5.92	6.85
p :	0.5	0.7	0.75	0.85	0.88	0.90

p represents the probability that the Z -test is better than the X -test. The same pattern is repeated if one makes the calculations for $N = 20$. Clearly the Z -test is on the whole a better test.

3.4 Corresponding test for the circle.

For the von Mises distribution also, $Z = R-X$ is the likelihood-ratio test statistic for H_0 ; the exact distribution cannot be handled, but the asymptotic result is $2\kappa(R-X) \approx \chi^2_1$, so that critical values z of Z would be given by

$$2\kappa z = \chi^2_1(\alpha) .$$

Monte Carlo results suggest that this approximation will hold with in a manner similar to that for the sphere, i.e., for $\kappa \geq 2$, and $N \geq 20$, and, for larger κ , even for $N \leq 20$.

The author gratefully acknowledges correspondence with Professor E.S. Pearson, on which Section 2 is almost entirely based.

This research has been supported by Contract No. N00014-67-A-0112-0053 (NR-042-267), Dept. of Statistics, Stanford University.

TABLE 2. Critical Values of R-X, for Test of Modal Vector, Sphere.

In descending order the values are for $\alpha = .01, .025, .05, .10$

N	k	0.5	1.0	1.5	2.0	2.5	3.0
4		4.722	3.746	2.902	2.273	1.837	1.534
		4.057	3.113	2.357	1.828	1.473	1.229
		3.486	2.597	1.931	1.487	1.197	0.998
		2.847	2.047	1.495	1.145	0.920	0.767
5		5.161	3.950	2.969	2.289	1.840	1.535
		4.407	3.258	2.399	1.837	1.475	1.229
		3.767	2.701	1.959	1.493	1.198	0.999
		3.06	2.117	1.512	1.149	0.921	0.767
6		5.532	4.099	3.008	2.296	1.841	1.535
		4.699	3.361	2.423	1.841	1.475	1.230
		4.000	2.775	1.974	1.496	1.198	0.999
		3.230	2.165	1.522	1.150	0.921	0.768
7		5.850	4.211	3.032	2.299	1.841	1.535
		4.948	3.438	2.437	1.843	1.475	1.230
		4.197	2.828	1.983	1.497	1.197	0.998
		3.376	2.200	1.527	1.150	0.920	0.766
8		6.127	4.296	3.046	2.300	1.841	1.535
		5.164	3.495	2.445	1.843	1.474	1.230
		4.307	2.868	1.988	1.497	1.197	0.999
		3.500	2.225	1.530	1.150	0.920	0.768
9		6.372	4.361	3.055	2.301	1.840	1.535
		5.354	3.538	2.450	1.843	1.474	1.230
		4.515	2.897	1.992	1.497	1.197	0.999
		3.608	2.243	1.532	1.151	0.920	0.768
10		6.590	4.412	3.060	2.301	1.839	1.535
		5.522	3.571	2.453	1.843	1.472	1.230
		4.645	2.919	1.994	1.496	1.198	0.999
		3.702	2.257	1.533	1.151	0.920	0.768
11		6.783	4.451	3.064	2.301	1.837	1.535
		5.669	3.596	2.455	1.843	1.472	1.230
		4.759	2.936	1.955	1.497	1.198	0.999
		3.784	2.267	1.534	1.151	0.920	0.765
12		6.956	4.482	3.065	2.299	1.837	1.535
		5.802	3.615	2.456	1.843	1.471	1.230
		4.862	2.949	1.994	1.497	1.198	0.999
		3.857	2.274	1.533	1.151	0.918	0.756
∞		9.210	4.605	3.070	2.302	1.842	1.535
		7.378	3.689	2.460	1.844	1.476	1.230
		5.991	2.996	1.997	1.498	1.198	0.999
		4.605	2.303	1.535	1.152	0.922	0.768

REFERENCES

- Stephens, M.A. (1962a). Exact and approximate tests for directions, 1. Biometrika, 49, 463-477.
- Stephens, M.A. (1962b). Exact and approximate tests for directions, 2. Biometrika, 49, 547-552.
- Stephens, M.A. (1967). Tests for the dispersion, and for the modal vector, of a distribution on a sphere. Biometrika, 54, 211-223.
- Stephens, M.A. (1969). Tests for the von Mises distribution. Biometrika, 56, 149-160.
- Watson, G.S. (1956). Analysis of dispersion on a sphere. Mon. Not. R. Astr. Soc. Geophys. Suppl. 7, 153-159.
- Watson, G.S. and Williams, E.J. (1956). On the construction of significance tests on the circle and the sphere. Biometrika, 43, 344-352.